Neuroscience Letters, 159 (1993) 236–238
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Abe, T., see Tohgi, H., 135
Agapito, M.T., see Pablos, M.I., 211
Al-Yousuf, S.A.H., Pow, D.V. and Mizuno, N., Ultrastructural demonstration of exocytosis of neurosecretory granules in the neurohypophysis of the frog *Rana temporalia*, 32
Ammendola, R., see Russo, T., 159
Arancibia, S., see Benyassi, A., 166
Askanas, V., Bilak, M., Engel, W.K., Leclerc, A. and Tomé, F., Prion protein is strongly immunolocalized at the postsynaptic domain of human normal neuromuscular junctions, 111
Astier, H., see Benyassi, A., 166

Badoer, E., Oldfield, B.J. and McKinley, M.J., Haemorrhage-induced production of Fos in neurons of the lamina terminalis: role of endogenous angiotensin II, 151
Beckh, S., see Swandulla, D., 59
Belin, M.-F., see Lepetit, P., 171
Benyassi, A., Roussel, J.-P., Rougeot, C., Gavaldà, A., Astier, H. and Arancibia, S., Chronic stress affects in vivo hypothalamic somatostatin release but not in vitro GH responsiveness to somatostatin in rats, 166
Berthelot, P., see Lacey, G., 64
Bilak, M., see Askanas, V., 111
Bobillier, P., see Lepetit, P., 171
Brecha, N.C., see Zittel, T.T., 79

Chai, C.-Y., see Chuang, S.-F., 127
Chan, J.Y.H., see Yien, H.-W., 9
Chan, S.H.H., see Yen, J.-C., 5
Chan, S.H.H., see Yien, H.-W., 9
Chiba, T., see Hanazawa, T., 71
Christofi, F.L., Hanani, M., Maudlej, N. and Wood, J.D., Enteric glial cells are major contributors to formation of cyclic AMP in myenteric plexus cultures from adult guinea-pig small intestine, 107
Chuang, S.-F., Chai, C.-Y. and Lin, M.-T., The interaction between locomotion, striatal dopamine and paramedian reticular nucleus in rats, 127
Cimino, F., see Russo, T., 159
Conejero, C., see Galarreta, M., 55
Connor, J.R., Tucker, P., Johnson, M. and Snyder, B., Ceruloplasmin levels in the human superior temporal gyrus in aging and Alzheimer's disease, 88
Córdova, M.D., see Pablos, M.I., 211
Couture, R., see Prat, A., 95
Curtis, D.R., see Lacey, G., 64

De Giorgio, R., see Zittel, T.T., 79
De Natale, G., see Desole, M.S., 143
Debaert, M., see Lacey, G., 64
del Río, R.M., see Galarreta, M., 55
Desole, M.S., Esposito, G., Enrico, P., Miele, M., Fresu, L., De Natale, G., Miele, E. and Grella, G., Effects of ageing on 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) neurotoxic effects on striatum and brainstem in the rat, 143
Devi, R.P.R. and Murthy, C.R.K., Ammonia-induced alterations in the activities of synaptosomal cholinesterases of rat brain under in vitro and in vivo conditions, 131
Doherty, P., see Harper, S.J., 202

Egger, C., Kirchmair, R., Hogue-Angeletti, R., Fischer-Colbrie, R. and Winkler, H., Different degrees of processing of secretogranin II in large dense core vesicles of bovine adrenal medulla and sympathetic axons correlate with their content of soluble PC1 and PC2, 199
Eifuku, S., see Ono, T., 75
Engel, W.K., see Askanas, V., 111
Enrico, P., see Desole, M.S., 143
Esposito, G., see Desole, M.S., 143

Facci, L., see Fadda, E., 147
Fadda, E., Negro, A., Facci, L. and Skaper, S.D., Ganglioside GM1 cooperates with brain-derived neurotrophic factor to protect dopaminergic neurons from 6-hydroxydopamine-induced degeneration, 147
Fatatis, A., see Russo, T., 159
Fevre-Montange, M., see Lepetit, P., 171
Fiore, F., see Russo, T., 159
Fischer-Colbrie, R., see Egger, C., 199
Flouquet, N., see Lacey, G., 64
Fresu, L., see Desole, M.S., 143
Fukata, J., see Nakaishi, S., 29
Fukushima, M., see Nakaishi, S., 29

Gähwiler, B.H., see Xie, X., 46
Galarreta, M., Solís, J.M., Menéndez, N., Conejero, C. and del Río, R.M., Nicotinamide adenine dinucleotides mimic adenosine inhibition on synaptic transmission by decreasing glutamate release in rat hippocampal slices, 55
Gavaldà, A., see Benyassi, A., 166
Gay, N., see Lepetit, P., 171
Gemba, H., Changes in cortical field potentials during learning processes of go/no-go reaction time hand movement with tone discrimination in the monkey, 21
Gerber, U., see Xie, X., 46
Gerilovsky, L., see Struppler, A., 115
Goshima, Y., see Yue, J.-L., 103
Grella, G., see Desole, M.S., 143
Grönroos, M. and Pertovaara, A., Capsaicin-induced central facilitation of a nociceptive flexion reflex in humans, 215

Hama, N., see Shirakami, G., 25
Hamato, H., see Tohgi, H., 135
Hanani, M., see Christofi, F.L., 107
Hanazawa, T., Motosugi, H., Konno, A., Kaneko, T., Tanaka, K. and Chiba, T., Distribution of NADPH-diaphorase positive nerve fibers in the rat nasal mucosa, 71
Harper, S.J., Walsh, F.S. and Doherty, P., Neurite outgrowth of spinal neurons on tissue sections of embryonic muscle is largely integrin dependent, 202
Hasegawa, H., see Wada, Y., 179
Hasséssian, H., see Prat, A., 95
Hershkowitz, N., see Katchman, A.N., 123
Hogue-Angeletti, R., see Egger, C., 199
Hoshino, M., Li, M., Zheng, L.Q., Suzuki, M., Mochizuki, T. and Yanaihara, N., Pituitary adenylate cyclase activating peptide and vasoactive intestinal polypeptide: differentiation effects on human neuroblastoma NB-OK-1 cells, 35
Huang, S.-K. and Pan, J.-T., Potentiating effects of serotonin and vasoactive intestinal peptide on the action of glutamate on suprachiasmatic neurons in brain slices, 1

Neuroscience Letters, 159 (1993) 236–238
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Abe, T., see Tohgi, H., 135
Agapito, M.T., see Pablos, M.I., 211
Al-Yousuf, S.A.H., Pow, D.V. and Mizuno, N., Ultrastructural demonstration of exocytosis of neurosecretory granules in the neurohypophysis of the frog *Rana temporalia*, 32
Ammendola, R., see Russo, T., 159
Arancibia, S., see Benyassi, A., 166
Askanas, V., Bilak, M., Engel, W.K., Leclerc, A. and Tomé, F., Prion protein is strongly immunolocalized at the postsynaptic domain of human normal neuromuscular junctions, 111
Astier, H., see Benyassi, A., 166

Badoer, E., Oldfield, B.J. and McKinley, M.J., Haemorrhage-induced production of Fos in neurons of the lamina terminalis: role of endogenous angiotensin II, 151
Beckh, S., see Swandulla, D., 59
Belin, M.-F., see Lepetit, P., 171
Benyassi, A., Roussel, J.-P., Rougeot, C., Gavaldà, A., Astier, H. and Arancibia, S., Chronic stress affects in vivo hypothalamic somatostatin release but not in vitro GH responsiveness to somatostatin in rats, 166
Berthelot, P., see Lacey, G., 64
Bilak, M., see Askanas, V., 111
Bobillier, P., see Lepetit, P., 171
Brecha, N.C., see Zittel, T.T., 79

Chai, C.-Y., see Chuang, S.-F., 127
Chan, J.Y.H., see Yien, H.-W., 9
Chan, S.H.H., see Yen, J.-C., 5
Chan, S.H.H., see Yien, H.-W., 9
Chiba, T., see Hanazawa, T., 71
Christofi, F.L., Hanani, M., Maudlej, N. and Wood, J.D., Enteric glial cells are major contributors to formation of cyclic AMP in myenteric plexus cultures from adult guinea-pig small intestine, 107
Chuang, S.-F., Chai, C.-Y. and Lin, M.-T., The interaction between locomotion, striatal dopamine and paramedian reticular nucleus in rats, 127
Cimino, F., see Russo, T., 159
Conejero, C., see Galarreta, M., 55
Connor, J.R., Tucker, P., Johnson, M. and Snyder, B., Ceruloplasmin levels in the human superior temporal gyrus in aging and Alzheimer's disease, 88
Córdova, M.D., see Pablos, M.I., 211
Couture, R., see Prat, A., 95
Curtis, D.R., see Lacey, G., 64

De Giorgio, R., see Zittel, T.T., 79
De Natale, G., see Desole, M.S., 143
Debaert, M., see Lacey, G., 64
del Río, R.M., see Galarreta, M., 55
Desole, M.S., Esposito, G., Enrico, P., Miele, M., Fresu, L., De Natale, G., Miele, E. and Grella, G., Effects of ageing on 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) neurotoxic effects on striatum and brainstem in the rat, 143
Devi, R.P.R. and Murthy, C.R.K., Ammonia-induced alterations in the activities of synaptosomal cholinesterases of rat brain under in vitro and in vivo conditions, 131
Doherty, P., see Harper, S.J., 202

Egger, C., Kirchmair, R., Hogue-Angeletti, R., Fischer-Colbrie, R. and Winkler, H., Different degrees of processing of secretogranin II in large dense core vesicles of bovine adrenal medulla and sympathetic axons correlate with their content of soluble PC1 and PC2, 199
Eifuku, S., see Ono, T., 75
Engel, W.K., see Askanas, V., 111
Enrico, P., see Desole, M.S., 143
Esposito, G., see Desole, M.S., 143

Facci, L., see Fadda, E., 147
Fadda, E., Negro, A., Facci, L. and Skaper, S.D., Ganglioside GM1 cooperates with brain-derived neurotrophic factor to protect dopaminergic neurons from 6-hydroxydopamine-induced degeneration, 147
Fatatis, A., see Russo, T., 159
Fevre-Montange, M., see Lepetit, P., 171
Fiore, F., see Russo, T., 159
Fischer-Colbrie, R., see Egger, C., 199
Flouquet, N., see Lacey, G., 64
Fresu, L., see Desole, M.S., 143
Fukata, J., see Nakaishi, S., 29
Fukushima, M., see Nakaishi, S., 29

Gähwiler, B.H., see Xie, X., 46
Galarreta, M., Solís, J.M., Menéndez, N., Conejero, C. and del Río, R.M., Nicotinamide adenine dinucleotides mimic adenosine inhibition on synaptic transmission by decreasing glutamate release in rat hippocampal slices, 55
Gavaldà, A., see Benyassi, A., 166
Gay, N., see Lepetit, P., 171
Gemba, H., Changes in cortical field potentials during learning processes of go/no-go reaction time hand movement with tone discrimination in the monkey, 21
Gerber, U., see Xie, X., 46
Gerilovsky, L., see Struppler, A., 115
Goshima, Y., see Yue, J.-L., 103
Grella, G., see Desole, M.S., 143
Grönroos, M. and Pertovaara, A., Capsaicin-induced central facilitation of a nociceptive flexion reflex in humans, 215

Hama, N., see Shirakami, G., 25
Hamato, H., see Tohgi, H., 135
Hanani, M., see Christofi, F.L., 107
Hanazawa, T., Motosugi, H., Konno, A., Kaneko, T., Tanaka, K. and Chiba, T., Distribution of NADPH-diaphorase positive nerve fibers in the rat nasal mucosa, 71
Harper, S.J., Walsh, F.S. and Doherty, P., Neurite outgrowth of spinal neurons on tissue sections of embryonic muscle is largely integrin dependent, 202
Hasegawa, H., see Wada, Y., 179
Hasséssian, H., see Prat, A., 95
Hershkowitz, N., see Katchman, A.N., 123
Hogue-Angeletti, R., see Egger, C., 199
Hoshino, M., Li, M., Zheng, L.Q., Suzuki, M., Mochizuki, T. and Yanaihara, N., Pituitary adenylate cyclase activating peptide and vasoactive intestinal polypeptide: differentiation effects on human neuroblastoma NB-OK-1 cells, 35
Huang, S.-K. and Pan, J.-T., Potentiating effects of serotonin and vasoactive intestinal peptide on the action of glutamate on suprachiasmatic neurons in brain slices, 1

Imura, H., see Nakaishi, S., 29
Inoue, O., see Suhara, T., 207
Itoh, H., see Shirakami, G., 25
Itoh, T., see Suhara, T., 207

Jakob, C., see Struppler, A., 115
Jiang, X., see Sublette, E., 175
Jingami, H., see Nakaishi, S., 29
Johnson, M., see Connor, J.R., 88

Kamiya, H.-o., see Saigo, A., 187
Kanda, M., see Song, W.-J., 99
Kanda, M., see Song, W.-J., 99
Kaneko, T., see Hanazawa, T., 71
Katchman, A.N. and Hershkowitz, N., Adenosine antagonists prevent hypoxia-induced depression of excitatory but not inhibitory synaptic currents, 123
Kayahara, T., see Yasui, Y., 39
Kirchmair, R., see Egger, C., 199
Kobayashi, K., see Suhara, T., 207
Komatsu, Y., see Shirakami, G., 25
Konno, A., see Hanazawa, T., 71
Kowall, N.W., see Nihei, K., 67
Krieger, C., Wagey, R. and Shaw, C., Amyotrophic lateral sclerosis: quantitative autoradiography of [^{3}H]MK- 801/NMDA binding sites in spinal cord, 191

Lacey, G., Berthelot, P., Vaccher, C., Flouquet, N., Vaccher, M.P., Debaert, M. and Curtis, D.R., Thienyl-GABA derivatives as specific baclofen agonists in the rat and cat spinal cord in vivo, 64
Leclerc, A., see Askanas, V., 111
Lee, T.-Y., see Yien, H.-W., 9
Lepetit, P., Fevre-Montange, M., Gay, N., Belin, M.-F. and Bobillier, P., Vasopressin mRNA in the cerebellum and circumventricular organs: a quantitative in situ hybridization study, 171
Li, M., see Hoshino, M., 35
Lin, M.-T., see Chuang, S.-F., 127
Lovinger, D.M., High ethanol sensitivity of recombinant AMPA-type glutamate receptors expressed in mammalian cells, 83

Martin, J.H., Differential spinal projections of subregions in the forelimb area of the motor cortex in the cat, 195
Matsumoto, T., see Saigo, A., 187
Maudlej, N., see Christofi, F.L., 107
McCabe, T. and Simon, R.P., Hyperthermia induces 72kDa heat shock protein expression in rat brain in non-neuronal cells, 163
McKinley, M.J., see Badoer, E., 151
Menéndez, N., see Galarreta, M., 55
Mesuraca, M., see Russo, T., 159
Miele, E., see Desole, M.S., 143
Miele, M., see Desole, M.S., 143
Misgeld, U., see Swandulla, D., 59
Misu, Y., see Yue, J.-L., 103
Mizuno, N., see Al-Yousuf, S.A.H., 32
Mochizuki, T., see Hoshino, M., 35
Mogavero, A.R., see Russo, T., 159
Mori, J.O., see Pablos, M.I., 211
Mori, K., see Shirakami, G., 25
Motosugi, H., see Hanazawa, T., 71
Murakami, F., see Song, W.-J., 99
Murthy, C.R.K., see Devi, R.P.R., 131

Naik, M.U., see Sublette, E., 175
Naito, Y., see Nakaishi, S., 29
Nakai, Y., see Nakaishi, S., 29
Nakaishi, S., Nakai, Y., Fukata, J., Naito, Y., Usui, T., Fukushima, M., Jingami, H., Nakao, K. and Imura, H., Immunoreactive corticotropin-releasing hormone levels in discrete hypothalamic nuclei of genetically obese Zucker rats, 29
Nakamura, K., see Ono, T., 75
Nakamura, M., see Wada, Y., 179
Nakano, K., see Yasui, Y., 39
Nakao, K., see Shirakami, G., 25
Nakao, K., see Nakaishi, S., 29
Nakayama, Y., see Saigo, A., 187
Negro, A., see Fadda, E., 147
Nihei, K. and Kowall, N.W., Involvement of NPY-immunoreactive neurons in the cerebral cortex of amyotrophic lateral sclerosis patients, 67
Nishijo, H., see Ono, T., 75

Ogawa, H., see Yamashita, Y., 155
Ohno, T., see Song, W.-J., 99
Ohno-Shosaku, T., Sawada, S. and Yamamoto, C., ATP-sensitive K^+ channel activators suppress the GABAergic inhibitory transmission by acting on both presynaptic and postsynaptic sites in rat cultured hippocampal neurons, 139
Ohnot, S., see Sublette, E., 175
Oikawa, H., see Song, W.-J., 99
Oldfield, B.J., see Badoer, E., 151
Ono, T., Eifuku, S., Nakamura, K. and Nishijo, H., Monkey hippocampal neuron responses related to spatial and non-spatial influence, 75
Osada, S.-i., see Sublette, E., 175
Osten, P., see Sublette, E., 175

Pablos, M.I., Agapito, M.T., Recio, J.M., Pérez-Gallardo, L., Córdova, M.D. and Mori, J.O., Effect of iron and estrogen on melatonin secretion by the chicken pineal gland, 211
Pan, J.-T., see Huang, S.-K., 1
Pardo, P.J. and Sams, M., Human auditory cortex responses to rising versus falling glides, 43
Pérez-Gallardo, L., see Pablos, M.I., 211
Pertovaara, A., see Grönroos, M., 215
Plenderleith, M.B. and Snow, P.J., The plant lectin *Bandeiraea simplicifolia* I-B_4 identifies a subpopulation of small diameter primary sensory neurones which innervate the skin in the rat, 17
Pow, D.V., see Al-Yousuf, S.A.H., 32
Prat, A., Hasséssian, H. and Couture, R., Neuropeptide K potently stimulates the hydrolysis of phosphatidylinositol in the rat spinal cord, 95
Puil, E., see Ries, C.R., 91

Raybould, H.E., see Zittel, T.T., 79
Recio, J.M., see Pablos, M.I., 211
Reuter, G., see Wicher, D., 13
Ries, C.R. and Puil, E., Isoflurane prevents transitions to tonic and burst firing modes in thalamic neurons, 91
Ríos, C., see Santamaría, A., 51
Rougeot, C., see Benyassi, A., 166
Roussel, J.-P., see Benyassi, A., 166
Russo, T., Mogavero, A.R., Ammendola, R., Mesuraca, M., Fiore, F., Fatatis, A., Salvatore, G. and Cimino, F., Immortalization of a cell

line showing some characteristics of the oligodendrocyte phenotype, 159

Sacktor, T.C., see Sublette, E., 175
Saigo, A., Takano, Y., Matsumoto, T., Tran, M., Nakayama, Y., Saito, R., Yamada, K. and Kamiya, H.-o., Central administration of senktide, a tachykinin NK-3 agonist, has an antidiuretic action by stimulating AVP release in water-loaded rats, 187
Saito, R., see Saigo, A., 187
Salvatore, G., see Russo, T., 159
Sams, M., see Pardo, P.J., 43
Santamaría, A. and Ríos, C., MK-801, an *N*-methyl-D-aspartate receptor antagonist, blocks quinolinic acid-induced lipid peroxidation in rat corpus striatum, 51
Sawada, S., see Ohno-Shosaku, T., 139
Shaw, C., see Krieger, C., 191
Shirakami, G., Itoh, H., Suga, S.-i., Komatsu, Y., Hama, N., Mori, K. and Nakao, K., Central action of C-type natriuretic peptide on vasopressin secretion in conscious rats, 25
Shiroyama, T., see Yasui, Y., 39
Simon, R.P., see McCabe, T., 163
Skaper, S.D., see Fadda, E., 147
Smart, T.G., see Xie, X., 46
Snow, P.J., see Plenderleith, M.B., 17
Snyder, B., see Connor, J.R., 88
Solís, J.M., see Galarreta, M., 55
Song, W.-J., Kanda, M., Ohno, T., Kanda, M., Oikawa, H. and Murakami, F., Segregation of cerebrorubral and cerebellorubral synaptic inputs on rubrospinal neurons of fetal cats as demonstrated by intracellular recording, 99
Sternini, C., see Zittel, T.T., 79
Struppler, A., Gerilovsky, L. and Jakob, C., Self-generated rapid taps directed to the opposite forearm in man: anticipatory reduction in the muscle activity of the target arm, 115
Sublette, E., Naik, M.U., Jiang, X., Osten, P., Valsamis, H., Osada, S.-i., Ohnot, S. and Sacktor, T.C., Evidence for a new, high-molecular weight isoform of protein kinase C in rat hippocampus, 175
Suga, S.-i., see Shirakami, G., 25
Suhara, T., Inoue, O., Kobayashi, K., Suzuki, K., Itoh, T. and Tateno, Y., No age-related changes in human benzodiazepine receptor binding measured by PET with [^{11}C]Ro 15-4513, 207
Suzuki, K., see Suhara, T., 207
Suzuki, M., see Hoshino, M., 35
Swandulla, D., Misgeld, U. and Beckh, S., Molecular properties of the glutamate receptor mediating synaptic excitation in rat hypothalamic neurons, 59

Takahashi, J., see Tohgi, H., 135
Takahashi, S., see Tohgi, H., 135
Takano, Y., see Saigo, A., 187
Tanaka, K., Cloning and expression of a glutamate transporter from mouse brain, 183
Tanaka, K., see Hanazawa, T., 71
Tateno, Y., see Suhara, T., 207
Tohgi, H., Abe, T., Takahashi, S., Takahashi, J. and Hamato, H., Alterations in the concentration of serotonergic and dopaminergic substances in the cerebrospinal fluid of patients with Parkinson's disease, and their changes after L-dopa administration, 135
Toida, K., see Yamashita, Y., 155
Tomé, F., see Askanas, V., 111
Tran, M., see Saigo, A., 187
Tsai, H.-F., see Yien, H.-W., 9
Tucker, P., see Connor, J.R., 88

Usui, T., see Nakaishi, S., 29

Vaccher, C., see Lacey, G., 64
Vaccher, M.P., see Lacey, G., 64
Valsamis, H., see Sublette, E., 175
Volonté, C., Dexamethasone abolishes the activation by nerve growth factor of protein kinase N: effects of nerve growth factor and dexamethasone on protein kinase N, 119

Wada, Y., Nakamura, M., Hasegawa, H. and Yamaguchi, N., Intrahippocampal injection of 8-hydroxy-2-(di-*n*-propylamino)tetralin (8-OH-DPAT) inhibits partial and generalized seizures induced by kindling stimulation in cats, 179
Wagey, R., see Krieger, C., 191
Walsh, F.S., see Harper, S.J., 202
Wicher, D. and Reuter, G., Neurohormone D increases the intracellular Ca^{2+} level in cockroach neurones through a Cd^{2+}-sensitive Ca^{2+} influx, 13
Winkler, H., see Egger, C., 199
Wood, J.D., see Christofi, F.L., 107

Xie, X., Gerber, U., Gähwiler, B.H. and Smart, T.G., Interaction of zinc with ionotropic and metabotropic glutamate receptors in rat hippocampal slices, 46

Yamada, K., see Saigo, A., 187
Yamaguchi, N., see Wada, Y., 179
Yamamoto, C., see Ohno-Shosaku, T., 139
Yamashita, Y., Toida, K. and Ogawa, H., Observation of Merkel cells with scanning electron microscopy, 155
Yanaihara, N., see Hoshino, M., 35
Yasui, Y., Kayahara, T., Shiroyama, T. and Nakano, K., Neurons in the intertrigeminal region of the rat send projection fibers to the superior colliculus, 39
Yen, J.-C. and Chan, S.H.H., Passive biophysical membrane properties of nucleus reticularis gigantocellularis neurons in brain slices from the rat, 5
Yien, H.-W., Chan, J.Y.H., Tsai, H.-F., Lee, T.-Y. and Chan, S.H.H., Participation of nucleus reticularis gigantocellularis in the antinociceptive effect of angiotensin III in the rat, 9
Yue, J.-L., Goshima, Y. and Misu, Y., Transmitter-like L-3,4-dihydroxyphenylalanine tonically functions to mediate vasodepressor control in the caudal ventrolateral medulla of rats, 103

Zheng, L.Q., see Hoshino, M., 35
Zittel, T.T., De Giorgio, R., Brecha, N.C., Sternini, C. and Raybould, H.E., Abdominal surgery induces c-fos expression in the nucleus of the solitary tract in the rat, 79